

New subgenera and a new species of the genus *Raphignathus* Dugès (Prostigmata, Raphignathidae), with taxonomic notes on the genus *Neoraphignathus* Smiley & Moser

Eid Muhammad Khan¹, Muhammad Kamran¹, Jawwad Hassan Mirza¹, Fahad Jaber Alatawi¹

¹ Acarology Research Laboratory, Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia
Corresponding author: Fahad Jaber Alatawi (falatawi@ksu.edu.sa)

Abstract

Four new subgenera in the genus *Raphignathus* Dugès are hereby proposed: *Raphignathus* (*Raphignathus*), **subgen. nov.**, *Raphignathus* (*Monoraphignathus*), **subgen. nov.**, *Raphignathus* (*Diraphignathus*), **subgen. nov.**, and *Raphignathus* (*Triraphignathus*), **subgen. nov.** These subgenera are diagnosed by the number of setae on the interscutal membrane of females. A new species, *R. (D.) neohecmatanaensis* **sp. nov.**, is described and illustrated based on females collected from *Ziziphus spina-christi* Mill. (Rhamnaceae). The taxonomic status of the monotypic genus *Neoraphignathus* Smiley & Moser and three species (*R. evidus*, *R. hsiufui*, and *R. johnstoni*) are discussed. A key to world species of the family Raphignathidae is given.

Key words: Acari, new combinations, predatory mite, Raphignathoidea, Saudi Arabia



Academic editor: Xiao-Feng Xue

Received: 10 May 2023

Accepted: 30 July 2023

Published: 23 August 2023

ZooBank: <https://zoobank.org/91E50336-528C-43DE-8266-42D12D0C7C1C>

Citation: Khan EM, Kamran M, Mirza JH, Alatawi FJ (2023) New subgenera and a new species of the genus *Raphignathus* Dugès (Prostigmata, Raphignathidae), with taxonomic notes on the genus *Neoraphignathus* Smiley & Moser. ZooKeys 1176: 165–180. <https://doi.org/10.3897/zookeys.1176.106224>

Copyright: © Eid Muhammad Khan et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Members of the family Raphignathidae Kramer (Prostigmata, Raphignathoidea) are active predators feeding on small arthropods (Meyer and Ueckermann 1989). They are mostly found in humus soil under dense bushes, leaf litter, lichens, and mosses and on a wide range of plants (Fan and Zhang 2005). Raphignathids have cervical peritremes and contiguous coxae and are divided into two genera: *Raphignathus* Dugès (76 species) and *Neoraphignathus* Smiley & Moser (one species, *N. howei* Smiley & Moser) (Smiley and Moser 1968; Khanjani et al. 2013). These genera are mainly differentiated by the presence and absence of dorsal shields, respectively. *Raphignathus* species have a worldwide distribution, whereas *Neoraphignathus* is only known from Louisiana, USA (Beron 2020).

The genus *Raphignathus* (type species: *R. ruberrimus* Dugès) was diagnosed as having three, or sometimes four, dorsal shields (Atyeo et al. 1961). While discussing the chaetotaxy of the superfamily Raphignathoidea, Atyeo et al. (1963) provided a general description of the genus *Raphignathus*. They considered the number of setae on prodorsal shields and on interscutal membrane as important taxonomic characteristics. In addition, the number of setae on the interscutal membrane was also used to differentiate among the species in recently published diagnostic keys (Nasrollahi et al. 2018; Pishehvar and Khanjani

2021). The validity of some *Raphignathus* species has been questionable due to ontogenetic development, and more females need to be collected and observed (Dönel and Doğan 2011). *Raphignathus broomicus* Podder was considered a species inquirenda due to uncertain and doubtful characters (Doğan and Erman 2019). Until now, most of published work on the family Raphignathidae has been regional and includes identification keys for China, India, Iran, and Turkey (Fan and Yin 2000; Doğan and Erman 2019; Pishehvar and Khanjani 2021).

In the present study, four new subgenera of *Raphignathus* are erected based on the prominent and consistent morphological character (number of setae on interscutal membrane). A new species, *R. (D.) neohecmatanaensis* sp. nov. is described and illustrated based on females. Some taxonomic notes on the monotypic genus, *Neoraphignathus* Smiley & Moser, and the identity of three species (*R. evidus*, *R. hsiufui*, and *R. johnstoni*) are concisely discussed. A diagnostic key to the world species is also provided.

Material and method

All published taxonomic literature on the family Raphignathidae was critically reviewed to confirm the validity of the species, subgeneric divisions and to prepare a diagnostic key of world species. The new raphignathoid species was collected by shaking foliage of *Ziziphus spina-christi* Mill. (Rhamnaceae) over a white sheet of paper; mite specimens were preserved in small vials containing 70% ethanol. The specimens were permanently mounted on glass slides in Hoyer's medium and identified under a phase-contrast microscope (BX51, Olympus, Tokyo, Japan). All measurements of the holotype specimen are given in micrometers (µm), followed by those of paratypes in the parenthesis. The terminology and abbreviations used in the description of the new species follow those of Kethley (1990) and Grandjean (1939, 1944, 1946). The holotypes and paratypes were deposited at the King Saud Museum of Arthropods (**KSMA**, Acarology section), Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia.

Results

Four new subgenera of the genus *Raphignathus* are proposed: *Raphignathus* (*Raphignathus*), subgen. nov., *Raphignathus* (*Monoraphignathus*), subgen. nov., *Raphignathus* (*Diraphignathus*), subgen. nov., and *Raphignathus* (*Triraphignathus*), subgen. nov., on the basis of the number of setae on the interscutal membrane, a prominent and consistent morphological character. A new species, *R. (D.) neohecmatanaensis* sp. nov., is described and illustrated based on adult females. Furthermore, taxonomic notes on the status of the monotypic genus, *Neoraphignathus* Smiley & Moser, and the identity of three species (*R. evidus*, *R. hsiufui* and *R. johnstoni*) are discussed. A diagnostic key to the world species is also presented.

Family Raphignathidae Kramer, 1877

Raphignathidae Kramer, 1877: 215

Type genus. *Raphignathus* Dugès, 1834: 53

Diagnosis (based on Krantz and Walter (2009) and Fan and Zhang (2005)). Peritremes linear, not imbedded in dorsal surface of stylophore; paired peritremes running laterally from base of stylophore to make short loops in collar membrane between gnathosoma and podosoma; coxae II and III contiguous; stigmata opening at base of chelicerae.

Taxonomic divisions of the genus *Raphignathus*

The presence or absence of prodorsal shields are diagnostic for the differentiation of the two existing raphignathid genera. Atyeo (1963) discussed in detail the chaetotaxy of the superfamily Raphignathoidea while describing seven *Raphignathus* species. This author also provided comprehensive diagnosis of *Raphignathus* and stated that the number of setae on the shields and the interscutal membrane is a consistent and important diagnostic character. The number of setae on the interscutal membrane is considered to be a strong character and has been used in identification keys to distinguish species (Nasrollahi et al. 2018; Pishehvar and Khanjani 2021). Through our extensive study of the literature of all 76 *Raphignathus* species, we find that the number of setae on the interscutal membrane can be used to erect subgenera.

In the current study, we categorize species of the genus *Raphignathus* into four new subgenera based on the number of setae on the interscutal membrane. These four subgenera are *Raphignathus* (*Raphignathus*), subgen. nov. without setae (11 spp.), *Raphignathus* (*Monoraphignathus*), subgen. nov. with one seta (14 spp.), *Raphignathus* (*Diraphignathus*), subgen. nov. with two setae (33 spp.), and *Raphignathus* (*Triraphignathus*), subgen. nov. with three setae (10 spp.).

***Raphignathus* (*Raphignathus*), subgen. nov.**

<https://zoobank.org/4A81EA35-DC89-4971-BD24-0459A27A8621>

Type species. *Raphignathus ruberrimus* Dugès, 1834: 53.

Diagnosis. Interscutal membrane without setae.

Etymology. The subgeneric epithet refers to the nominotypical subgenus.

***Raphignathus* (*Monoraphignathus*), subgen. nov.**

<https://zoobank.org/D332A8E8-0390-4D39-95D4-A8C6E7E0B8D5>

Type species. *Raphignathus bathursti* Meyer & Ryke, 1960: 229.

Diagnosis. Interscutal membrane with one pair of setae.

Etymology. The subgeneric epithet refers to the one pair of setae on interscutal membrane.

***Raphignathus* (*Diraphignathus*), subgen. nov.**

<https://zoobank.org/0D7C8B12-10E8-4019-A2FA-591934FD9B17>

Type species. *Raphignathus gracilis* (Rack, 1962): 281.

Diagnosis. Interscutal membrane with two pairs of setae.

Etymology. The sub-generic epithet refers to the two pairs of setae on interscutal membrane.

The species included in this new subgenus are widely distributed over the world.

***Raphignathus* (*Triraphignathus*), subgen. nov.**

<https://zoobank.org/80C632B9-6521-47BB-A6E2-3EA44EE761DD>

Type species. *Raphignathus domesticus* Shiba, 1969: 157.

Diagnosis. Interscutal membrane with three pairs of setae.

Etymology. The subgeneric epithet refers to the interscutal membrane with three pairs of setae.

Notes on the taxonomic status of the genus *Neoraphignathus*

To date, the family Raphignathidae has included two genera, *Raphignathus* and *Neoraphignathus*, which have been differentiated based on the presence or absence of shields on the dorsum. The monotypic genus, *Neoraphignathus* (type species: *N. howei* Smiley & Moser) was erected in 1968, based on a single female holotype specimen without detailed description and illustration. Based on observations and the collection of the immature specimens of the genus *Raphignathus*, prodorsal shields are weakly developed or absent in immatures. Atyeo et al. (1961) has reported that dorsal shields are sometime feebly developed. Since its first description, the type species, *N. howei*, has not been redescribed, nor have new *Neoraphignathus* species been described. We suggest that the type specimen of *N. howei* be re-examined and also that more specimens be collected from the type locality to confirm the absence of a dorsal shield to confirm the validity of *Neoraphignathus*.

Notes on the validity of *Raphignathus evidus*, *R. hsiufui*, and *R. johnstoni*

The taxonomic identity of *Raphignathus evidus* Fan, *R. hsiufui* Fan, and *R. johnstoni* Womersley are doubtful. These species were originally described based on single specimens, minor differential characteristics (i.e. number of dorsal setae on the lateral prodorsal shield; all three species have two pairs of setae on lateral shields), and small opisthosomal shields. In contrast, all other *Raphignathus* species have three pairs of setae on the lateral prodorsal shields along with the pores (*ia*). The immature stages of *Raphignathus* gradually develop the prodorsal shields, striation patterns, and leg setae (Fan and Yin 2000). For instance, the immature stages of *R. giselae*, *R. lenis*, and *R. caspicus* each have two setae on the lateral prodorsal shields (three setae in adult) and small lateral prodorsal shields with weakly developed striations. Moreover, we also observed the immatures from more than 10 populations of *Raphignathus* and found reduced size of weakly sclerotized lateral prodorsal shields and setae set on the edges of shields. Based on this evidence, *R. evidus*, *R. hsiufui*, and *R. johnstoni* should be revised and more specimens collected to confirm their validity.

New species

Raphignathus (*Diraphignathus*) *neohecmatanaensis* sp. nov.

<https://zoobank.org/12E13CEE-478F-4A3F-A47E-B3C7F34C73FA>

Figs 1–8

Diagnosis. Female: endopodal shield absent between coxae I–IV; two small shields present posterolateral to median prodorsal shield; palp femora with two setae; femora 6–5–3–2; genua 5(+κ)–5(+κ)–4–4; tibiae 5(+φρ)–5(+φρ)–4(+φρ)–4(+φρ); tarsi 21(1ω)–20(1ω)–15–14.

Description. Female ($n = 4$). Idiosoma oval, length of body (including gnathosoma) 533 (525–545); width of body 345 (338–353).

Dorsum (Fig. 1). Propodosoma with one medial and two lateral shields each containing three setae; medial sclerite with setae *vi*, *si* and *c1*; paired ovoid lateral shields each with an eye, one cupule (*ia*) and *sci*, *sce* and seta *c2* seta; opisthosomal setal pairs *d1*, *e1* and the pair of cupuli (*im*); posterior opisthosomal shield large, rectangular, bearing four pairs of setae (*f1*, *h1*–3) and one pair of the cupule (*ip*); all dorsal shields finely punctate; dorsal body setae setiform, smooth and acute; pseudanal setae *ps1* dorsally located. Lengths of dorsal setae: *vi* 25 (23–27), *ve* 28 (26–30), *sci* 27 (26–29), *sce* 28 (26–29), *c1* 23 (17–19), *c2* 25 (19–23), *d1* 21 (20–23), *e1* 22 (21–23), *f1* 21 (19–22), *h1* 24 (23–25), *h2* 25 (23–25), *h3* 22 (21–24), *ps1* 23 (21–25); distances between dorsal 195 setae: *vi*–*vi* 27 (29–31), *sci*–*sci* 123 (118–128), *vi*–*sci* 70 (66–72), *sce*–*c2* 68 (65–72), *c1*–*c1* 30 (28–32), *d1*–*d1* 99 (92–101), *c1*–*d1* 53 (49–55), *d1*–*e1* 22 (21–24), *f1*–*f1* 88 (83–90), *e1*–*f1* 65 (61–68), *h1*–*h1* 45 (42–48), *h1*–*h2* 43 (41–46), *h2*–*h2* 77 (72–80), *h3*–*h3* 101 (98–106).

Venter (Fig. 2). Venter entirely striated, without punctations; coxisternal shields absent (Fig. 2); ventral setae lengths: *1a* 42 (41–44), *1b* 38 (35–39), *1c* 34 (33–36), *2b* 36 (30–34), *2c* 32 (30–34), *3a* 24 (23–26), *3b* 18 (17–20), *3c* 36 (34–38), *4a* 22 (23–27), *4c* 38 (35–40); two pairs of aggenital setae (*ag1*–2) with one cupule (*ih*) on each side of the genital shield; anal opening and genital shields separate; genital shield prominent with a few punctations, bearing three pairs of genital setae (*g1*–3); anal opening terminal, with three pairs of setae (*ps1*–3), *ps1* dorsal *ps3* and *ps2* ventral; ventral setal lengths: *ag1* 29 (27–32); *ag2* 27 (25–28); *g1* 31 (28–33); *g2* 25 (23–28); *g3* 20 (19–22); *ps2* 22 (21–23); *ps3* 21 (20–22). Distances between ventral setae: *1a*–*1a* 55 (53–58), *3a*–*3a* 115 (97–109), *4a*–*4a* 70 (68–73), *ag2*–*ag2* 42 (41–43), *g1*–*g1* 38 (35–40), *g2*–*g2* 52 (48–55), *g3*–*g3* 75 (74–75), *2b*–*2c* 30 (25–29), *1a*–*3a* 50 (49–53), *3a*–*4a* 72 (68–75), *4a*–*ag1* 85 (82–88), *ag1*–*ag2* 65 (63–68), *ag2*–*g1* 63 (60–65), *g1*–*g2* 15 (14–17), *g2*–*g3* 22 (21–25), *ag*–*g1* 42 (40–45), *g3*–*ps3* 16 (15–18), *ps2*–*ps3* 16 (14–18).

Gnathosoma (Figs 3, 4). Ventral infracapitular with two pairs of very long setae (*m* and *n*), *m* 40 (39–42), *n* 52 (49–54) and two pairs of pilose adoral setae (*or1*–2), *or1* 23 (21–24), *or2* 20 (19–22) (Fig. 3); stylophore conical and striated; palp chaetotaxy (femur-tarsus) as follows: 3–2–4+1 claw 4+1ω+4 eupathidia (ζ) (Fig. 4).

Legs (Figs 5–8). Length of legs I–IV (without coxae): 340 (328–348); 270 (276–286); 325 (317–328); 375 (367–384), respectively. Chaetotaxy on legs




Figure 1. *Raphignathus* (*Diraphignathus*) *neohecmatanaensis* sp. nov. (female), dorsum. Scale bar: 100 μ m.




Figure 2. *Raphignathus* (*Diraphignathus*) *neohecmatanaensis* sp. nov. (female), venter. Scale bar: 100 μ m.




Figure 3. *Raphignathus* (*Diraphignathus*) *neohecmatanaensis* sp. nov. (female), gnathosoma. Scale bar: 50 μ m.





Figure 4. *Raphignathus* (*Diraphignathus*) *neohecmatanaensis* sp. nov. (female), palp. Scale bar: 20 μ m.



Figures 5–8. *Raphignathus* (*Diraphignathus*) *neohecmatanaensis* sp. nov. (female), Legs I–IV. Scale bar: 20 μ m.

I–IV (solenidia in parentheses and not included in setal counts): coxa 2–2–2–1; trochanter 1–1–2–1; femora 6–5–3–2; genua 5(+ κ)-5(+ κ)-4–4; tibiae 5(+ $\varphi\rho$)-5(+ $\varphi\rho$)-4(+ $\varphi\rho$)-4(+ $\varphi\rho$); tarsi 21(1 ω)-20(1 ω)-15–14.

Male and immature stages. Unknown.

Type materials. *Holotype* female and three paratype females, Faifa, Jazan, 24°30.412'N, 39°36.578'E, 8 Oct., 2020, collected from *Ziziphus spina-christi* Mill. (Rhamnaceae) by Eid M. Khan, Jawwad H. Mirza & Hafiz S. Mushtaq.

Etymology. The specific epithet is in reference to the similarity of the new species to *R. (D.) hecmatanaensis*; *neo* = new.

Remarks. *Raphignathus (D.) neohecmatanaensis* sp. nov., belongs to the subgenus *Diraphignathus* subgen. nov. The new species resembles to *R. (D.) hecmatanaensis* Khanjani & Ueckermann in having two pairs of setae (*d1* and *e1*) on the interscutal membrane, two setae on the palp femora, and two small plates present posterolateral to the median prodorsal shield. However, the new species differs from *R. (D.) hecmatanaensis* in the absence of an endopodal shield (vs present), femur IV with two setae (vs three), and leg tarsus I with one solenidion (vs two solenidia).

Key to genera, subgenera, and all known species of the family Raphignathidae

Five species are not included in the key. *Raphignathus longimanus* (Koch), *R. impressus* (Koch), *R. hispidus* (Dugès), and *R. deserticula* (Trägårdh) because their descriptions are incomplete, and *R. lanuginosus* Atyeo is excluded, as it was described on the male.

- 1 Dorsum with well-developed shields, one medial, one pair of lateral shields on prodorsum and one hysterosomal shield..... **Genus *Raphignathus* [2]**
- Dorsum without shields.... **Genus *Neoraphignathus*, *N. howei* Smiley & Moser**
- 2 Interscutal membrane without setae, hysterosomal shield with six pairs of setae..... ***Raphignathus* (*Raphignathus*) subgen. nov. [3]**
- Interscutal membrane with ≥ 1 setae **13**
- 3 Opisthosoma or opisthosomal shield reticulated **4**
- Opisthosomal shield smooth or otherwise, not reticulated..... **6**
- 4 Prodorsal shields reticulated.... ***R. (R.) crustus* Fan & Zhang, New Zealand**
- Prodorsal shields smooth or punctate..... **5**
- 5 Tibia I with 5 +2 ϕ , Tibia III with 5+1 ϕ , dorsal body setae comparatively long (54–74) ***R. (R.) kurdistanensis* Nasrollahi, Khanjani & Mirfakhraei, Iran**
- Tibia I with 5 +1 ϕ , Tibia III with 4+1 ϕ , dorsal body setae comparatively short (24–36) ***R. (R.) darjeelingensis* Gupta, India**
- 6 Opisthosoma without distinct shield; setae *d1*, *e1*, and *f1* very minute, 1/3–1/2 length of setae *v1* ***R. (R.) guajavae* Gupta, India**
- Opisthosoma with distinct opisthosomal shield; setae *d1*, *e1*, and *f1* at least 2/3 length of setae *v1* **7**
- 7 Genu I and II each with a large leaf-like solenidion ***R. (R.) pycnonotus* (Gupta & Paul), India**
- Genu I and II each with a small, slender solenidion **8**
- 8 Setae *c1* short, reaching 1/3 length of interscutal membrane, far behind the anterior margin of opisthosomal shield; tarsus IV with large solenidion..... ***R. (R.) inornata* (Gupta & Paul), India**
- Setae *c1* crossing interscutal membrane and anterior margin of opisthosomal shield; tarsus IV with small solenidion **9**
- 9 Tibia I with one solenidion **10**
- Tibia I with two solenidia **12**
- 10 Genu IV with three setae..... ***R. (R.) hirtellus* Athias-Henriot, Algeria**
- Genu IV with four setae..... **11**
- 11 Ratios $d1-d1/c1-c1 = 3.00$, $c2-c2/d1-d1 = 3$, $d1-d1/e1-e1 = 0.58$, $e1-e1/f1-f1 = 0.94-1.00$ ***R. (R.) neocardinalis* Atyeo, The Bahamas**
- Ratios $d1-d1/c1-c1 = 6.00$, $c2-c2/d1-d1 = 1.22$, $d1-d1/e1-e1 = 1.38$, $e1-e1/f1-f1 = 0.65$ ***R. (R.) conspicuus* (Berlese), Colombia**
- 12 Dorsal setae comparatively long; setae *c1*, and *d1* crossing bases of *d1* and *e1*, respectively ***R. (R.) khorramabadensis* Bagheri, Jafari & Paktinat, Iran**
- Dorsal setae comparatively short; setae *c1*, and *d1* far behind bases of *d1* and *e1*, respectively ***R. (R.) cardinalis* (Ewing), USA**

- 13 Interscutal membrane with one pair of setae.....
.....***Raphignathus (Monoraphignathus)* subgen. nov. [14]**
- Interscutal membrane with more than one pair of setae27
- 14 Palp femur with two setae; femur I with three setae
.....***R. (M.) arabicus* Gomaa & Hassan, Egypt**
- Palp femur with three setae; femur I with five or six setae.....15
- 15 Femur IV with two or three setae16
- Femur IV with four setae.....21
- 16 Femur IV with two setae17
- Femur IV with three setae19
- 17 Genu II with five setae including micro setae
.....***R. (M.) costatus* Chaudhri, Akbar & Rasool, Pakistan**
- Genu II with six setae including microsetae18
- 18 Setae *e1* reaching to bases of *h1*; dorsal body setae with spinules along
entire length.....***R. (M.) zhaoi* Hu, Jing & Liang, China**
- Setae *e1* reaching half distance to bases of *h1* (or distance *e1–h1*), dorsal
body setae with spinules along entire length
.....***R. (M.) kuznetzovi* Dogan & Ayyildiz, Turkey**
- 19 Setae *c2* crossing bases of *d1*, Setae *c1* extending to bases of *e1*, setae *e1*
extending to bases of *h1****R. (M.) ueckermanni* Koç & Kara, Turkey**
- Setae *c2* crossing setae *c1* far behind to the bases of *e1*, setae *e1* far be-
hind to the bases of *h1*20
- 20 Dorsal body setae ensiform, setae *c1* far behind bases of *d1*, *d1–d1* dis-
tance almost five times more than *c1–c1* distance.....
.....***R. (M.) ensipilosus* Meyer & Ueckermann, South Africa**
- Dorsal body setae setiform, setae *c1* far behind bases of *d1*, *d1–d1* dis-
tance almost equal to *c1–c1****R. (M.) cometes* Atyeo, Bahama-Islands**
- 21 Small shields absent posterolateral to median prodorsal shield.....22
- Small shields present posterolateral to median prodorsal shield.....25
- 22 Genital plates/covers with four pairs of setae
.....***R. (M.) koseiensis* Dönel & Doğan, Turkey**
- Genital plates/covers with three pairs of setae23
- 23 Femur I and II each with five setae... ***R. (M.) solimani* Hassan & Gomaa, Egypt**
- Femur I and II each with six setae.....24
- 24 Dorsal setae comparatively long; most setae cross base of next consecu-
tive setae.....***R. (M.) kelkitensis* Dönel & Doğan, Turkey**
- Dorsal setae comparatively short; most setae far behind base of next con-
secutive setae.....***R. (M.) fani* Doğan & Ayyildiz, Turkey**
- 25 Dorsal setae comparatively long; most setae reach or cross base of next
consecutive setae***R. (M.) bathursti* Meyer & Ryke, South Africa**
- Dorsal setae comparatively short; most setae far behind base of next con-
secutive setae.....26
- 26 Trochanter III with three setae..... ***R. (M.) afyonensis* Akyol & Koç, Turkey**
- Trochanter III with two setae
.....***R. (M.) collegiatus* Atyeo, Baker, & Crossley, USA**
- 27 Interscutal membrane with two pairs of setae.....
.....***R. (Diraphignathus)* subgen. nov. [28]**
- Interscutal membrane with three or four pairs of setae
.....***R. (Triraphignathus)* subgen. nov. [60]**

28	Medial prodorsal shield with two pairs of setae	29
–	Medial prodorsal shield with three pairs of setae	30
29	Setae <i>c1</i> present, setae <i>vi</i> absents; presence of plates behind the anteromedian plate; femur IV with 2 setae	
 <i>R. (D.) evansi</i> Zaher & Gomaa, Egypt	
–	Setae <i>c1</i> absent, setae <i>vi</i> present; dorsum without a pair of small plates behind anteromedian plate; femur IV with 3 setae	
 <i>R. (D.) ehari</i> Zaher & Gomaa, Egypt	
30	Genital plates/covers with four pairs of setae	31
–	Genital plates/covers with three pairs of setae	32
31	Two small shields posterior to median prodorsal shield absent; endopodal shields absent; setae <i>f1</i> behind the anterior margin of opisthosomal shield.....	<i>R. (D.) saboorii</i> Ghorbani & Bagheri, Iran
–	Two small shields posterior to median prodorsal shield present; endopodal shields present; setae <i>f1</i> on the anterior margin of opisthosomal shield.....	<i>R. (D.) karabagiensis</i> Akyol & Koç, Turkey
32	Palp femur with two setae	33
–	Palp femur with three setae.....	44
33	Two small shields posterior to median prodorsal shield present	34
–	Two small shields posterior to median prodorsal shield absent	38
34	Endopodal shield present; femur IV with three setae	35
–	Endopodal shield absent; femur IV with two setae.....	36
35	Tarsus I with two solenidia	
 <i>R. (D.) hecmatanaensis</i> Khanjani & Ueckermann, Iran	
–	Tarsus I with one solenidion.....	<i>R. (D.) arcus</i> Akyol, Turkey
36	Femur IV with two setae; tarsus I with one solenidion	
 <i>R. (D.) neohecmatanaensis</i> sp. nov. Alatawi & Kamran, Saudi Arabia	
–	Femur IV with three setae; tarsus I with two solenidia	37
37	Lateral prodorsal shield with one pair of pob; tibiae III 5(+1 ϕ p) tarsi 18(+1 ω +1 ω 2)	<i>R. (D.) seraji</i> Pishehvar & Khanjani, Iran
–	Lateral prodorsal shield without pob; tibiae III 5 tarsi 19(+1 ω +1 ω 2)	
 <i>R. (D.) rakhshandehi</i> Pishehvar & Khanjani, Iran	
38	Dorsal setae distally forked or tricarinate.....	
 <i>R. (D.) furcisetosus</i> Meyer & Ueckermann, South Africa	
–	Dorsal setae simple, not distally forked or tricarinate	39
39	Femur IV with two setae	<i>R. (D.) erzincanica</i> Doğan, Turkey
–	Femur IV with three setae	40
40	Opisthosomal shield reduced; interscutal membrane more longer than opisthosomal shield.....	41
–	Opisthosomal shield equally long or longer than interscutal membrane	43
41	Dorsal setae stout, serrate and blunt-tipped	
 <i>R. (D.) membranus</i> Fan & Yin, China	
–	Dorsal setae simple, distally pointed	42
42	Interscutal membrane four times longer than much reduced opisthosomal shield; <i>f1</i> on anterior margin of opisthosomal shield	
 <i>R. (D.) vahiti</i> Doğan, Turkey	
–	Interscutal membrane slightly longer than opisthosomal shield; <i>f1</i> behind anterior margin of opisthosomal shield.....	
 <i>R. (D.) giselae</i> Meyer & Ueckermann, Zimbabwe	

- 43 Median prodorsal shield anteriorly extending to peritremes and wider anteriorly near setae *sci* as compared to posterior half; setae *f1* on anterior margin of opisthosomal shield.....***R. (D.) gracilis* (Rack), Germany**
 - Median prodorsal shield anteriorly far behind peritremes and almost equally wide anteriorly near setae *sci* and at posterior half; setae *f1* just behind anterior margin of opisthosomal shield***R. (D.) bakeri* Zaher & Gomaa, Egypt**
- 44 Small shields posterior to median prodorsal shield absent49
 - Two small shields posterior to median prodorsal shield present45
- 45 Coxae II with one seta ***R. (D.) atyeoi* Meyer & Ueckermann, South Africa**
 - Coxae II with two setae46
- 46 Femur IV with two setae47
 - Femur IV with three setae48
- 47 Coxae III and IV with endopodal shields; setae *f1* on posterior margin of interscutal membrane; distance $f1-f1 < d1-d1$***R. (D.) summersi* Robaux, USA**
 - Coxae III and IV without endopodal shields; setae *f1* far behind posterior margin of interscutal membrane; distance $f1-f1 > d1-d1$
.....***R. (D.) aciculatus* Fan, China**
- 48 Tarsus I–IV 19+ ω , 15+1 ω , 13, 12
.....***R. (D.) africanus* Meyer & Ueckermann, South Africa**
 - Tarsus I–IV 21+1 ω , 21+1 ω , 15, 14.....
.....***R. (D.) hatamii* Khanjani & Pishehvar, Iran**
- 49 Coxa II with one seta.....***R. (D.) rarus* Kuznetsov, USSR**
 - Coxa II with two setae50
- 50 Coxae III and IV without endopodal shields51
 - Coxae III and IV with endopodal shields.....53
- 51 $c1-f1/f1-f1 = 0.70$; $c1-f1 < f1-f1$; space between setae *f1-f1* twice as wide as between setae *d1-d1****R. (D.) atomatus* Fan & Zhang, New Zealand**
 - $c1-f1/f1-f1 = 1.50-1.87$; $c1-f1$ 1.5–2.0 times more than *f1-f1*; distance $f1-f1 \leq d1-d1$ 52
- 52 Dorsal setae barbed; setae *c1*, *d1*, and *e1* reach or cross bases of next consecutive setae, distances $d1-f1 \leq f1-f1$; setae *f1* near anterior margin of opisthosomal shield***R. (D.) satoi* Shiba, Malay Peninsula**
 - Dorsal setae simple; setae *c1*, *d1*, and *e1* not reaching base of next consecutive seta; distances $d1-f1$ 1.31 times more as *f1-f1*, *f1* behind anterior margin of opisthosomal shield
.....***R. (D.) kamiensis* Meyer & Ueckermann, South Africa**
- 53 Femur I and II with five and four setae, respectively
.....***R. (D.) hexeris* Chaudhri, Akbar & Rasool, Pakistan**
 - Femur I and II with six and five setae, respectively54
- 54 Opisthosomal shield 2–4 times wider than interscutal membrane.....55
 - Interscutal membrane as wide as or more wider than opisthosomal shield.....56
- 55 Setae *c1* far behind posterior margin of prodorsal shield; tibia III with five setae excluding solenidion; opisthosomal shield four times wider than interscutal membrane ***R. (D.) neogracilis* Robaux, USA**
 - Setae *c1* on the posterior margin of prodorsal shield; tibia III with four setae excluding solenidion; opisthosomal shield twice as wide as interscutal membrane..... ***R. (D.) scutatus* Kuznetsov, USSR**

56	Femur IV with two setae	57
–	Femur IV with three setae	59
57	Dorsal shields without striations, tarsus I with 22 setae	<i>R. (D.) tumidus</i> Kuznetsov, USSR
–	Dorsal shields with fine, sparse puncta and faint striae; tarsi I with 21 setae	58
58	Tarsi III–IV with 14 and 13 setae, respectively; femur II with five setae	<i>R. (D.) caspicus</i> Doustaresharaf and Kazemi, Colombia
–	Tarsi III–IV with 15 and 14 setae, respectively; femur II with six setae	<i>R. (D.) tamaricis</i> Poudineh, Ramroodi & Bagheri, Iran
59	Setae $f1-f1 \leq c1-c1$ and $d1-d1$	<i>R. (D.) giresuniensis</i> Doğan, Turkey
–	Setae $f1$ twice as widely spaced as $c1-c1$	<i>R. (D.) orientalis</i> Fan & Li, China
60	Medial prodorsal shield with two pairs of setae	<i>R. (T.) lenis</i> Barillo, Uzbekistan
–	Medial prodorsal shield with ≥ 3 pairs of setae	61
61	Genital shield with four pairs of setae	62
–	Genital shield with three pairs of setae	63
62	Two small shields present posterolateral to prodorsal shield; endopodal shields near coxae III and IV absent	<i>R. (T.) sceptrum</i> Chaudhri, Akbar & Rasool, Pakistan
–	Small shields absent posterolateral to prodorsal shield; endopodal shields near coxae III and IV present	<i>R. (T.) quadrigeminus</i> Dönel & Doğan, Turkey
63	Palp femur with three setae	<i>R. (T.) aethiopicus</i> (Meyer & Ryke), South Africa
–	Palp femur with two setae	64
64	Femur IV with two setae	<i>R. (T.) karrooi</i> Meyer & Ueckermann, South Africa
–	Femur IV with three setae	65
65	Femur I with five setae, femur II with four setae	<i>R. (T.) domesticus</i> Shiba, Japan
–	Femur I with six setae, femur II with five setae	66
66	Two small shields present posterolateral to prodorsal shield	67
–	Small shields absent posterolateral to prodorsal shield	68
67	Tibiae III with four setae, solenidia absent	<i>R. (T.) hamooniensis</i> Poudineh, Ramroodi & Bagheri, Iran
–	Tibiae III with five setae with one solenidion	<i>R. (T.) larestanensis</i> Bagheri, Akrami & Majidi, Iran
68	Genu II with four tactile setae; endopodal shields near coxae I–II present	<i>R. (T.) emirdagiensis</i> Akyol & Koç, Turkey
–	Genu II with five tactile setae; endopodal shields near coxae I–II absent	<i>R. (T.) ozkani</i> Doğan, Turkey

Discussion

The taxonomic classification of predatory mites of the genus *Raphignathus* are revised, and for the first time, the genus is divided into four subgenera by con-

sidering the morphologically valid, persistent, and prominent characters (Atyeo 1963). The use of subgenera supports the identification of raphignathoid species and will help to avoid designation of new species based on variable characters. *Raphignathus evidus*, *R. hsiufui*, and *R. johnstoni* are considered doubtfully valid. They were described based on size of lateral prodorsal shields and number of setae, but these in these characteristics they resemble immature stages (Fan and Yin 2000). The monotypic genus *Neoraphignathus* was erected based on only the single type specimen with a restricted geographical region and its description is brief. We suspect it might have been described based on the immature stage of a *Raphignathus* species, and we suggest that the type species be revised and more specimens collected to confirm the validity of the species and genus.

Acknowledgements

The authors extend their appreciation and special thanks to Edward Ueckermann (North-West University, South Africa), Carlos Holger Wenzel Flechtmann (Universidade de São Paulo, Piracicaba, São Paulo, Brazil), Zhi-Qiang Zhang (Landcare Research, Auckland, New Zealand), and Daochao Jin (Guizhou University, Guiyang, China) for providing useful literature.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The authors would like to extend their sincere appreciation to the researchers supporting project number (RSPD2023R807), King Saud University, Riyadh, Saudi Arabia.

Author contributions

Conceptualization: MK, FJJA. Data curation: JHHM, EMK. Funding acquisition: FJJA. Investigation: EMK, MK, FJJA. Methodology: JHHM, EMK. Resources: FJJA. Supervision: FJJA. Visualization: JHHM, EMK. Writing - original draft: JHHM, EMK. Writing - review and editing: FJJA, MK.

Author ORCIDs

Eid Muhammad Khan  <https://orcid.org/0000-0001-5008-2524>

Muhammad Kamran  <http://orcid.org/0000-0001-6084-203X>

Jawwad Hassan Mirza  <https://orcid.org/0000-0002-1726-4331>

Fahad Jaber Alatawi  <https://orcid.org/0000-0002-6824-2650>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Atyeo WT (1963) New and re-described species of Raphignathidae (Acarina) and discussion of the chaetotaxy of Raphignathoidea. *Journal of the Kansas Entomological Society* 36: 172–186.
- Atyeo WT, Baker EA, Crossley DA (1961) The genus *Raphignathus* Dugés (Acarina: Raphignathoidea) in the United State with notes on the Old World species. *Acarologia* 3(1): 14–20.
- Beron P (2020) *Acarorum catalogus VII: Trombidiformes, Prostigmata, Raphignathoidea* (Fam. Barbutiidae, Caligonellidae, Camerobiidae, Cryptognathidae, Dasythyreidae, Dytiscacaridae, Eupalopsellidae, Homocaligidae, Mecognathidae, Raphignathidae, Stigmaeidae, Xenocaligonellidae). Pensoft – National Museum of Natural History – Bulgarian Academy of Sciences, Sofia, 306 pp. <https://doi.org/10.3897/ab.e55087>
- Doğan S, Erman O (2019) Mites of *Raphignathus* Dugés (Acari: Raphignathidae) from Harşit Valley (Turkey), with taxonomic notes on other members of the genus. *Bitki Koruma Bulteni* 59(1): 25–36. <https://doi.org/10.16955/bitkorb.419431>
- Dönel G, Doğan S (2011) A new species of *Raphignathus* Dugés (Acari, Raphignathidae) and newly discovered male of *R. fani* Doğan and Ayyıldız from Turkey. *International Journal of Acarology* 37(sup1, supplement 1): 27–33. <https://doi.org/10.1080/01647954.2010.531765>
- Dugès A (1834) Sur les acariens. *Annales des Science Naturelle* 29. Série Zoologie 2: 5–46.
- Fan QH, Yin XM (2000) The genus *Raphignathus* (Acari: Raphignathidae) from China. *Systematic and Applied Acarology* 5(0): 83–98. <https://doi.org/10.11158/saa.5.1.11>
- Fan QH, Zhang ZQ (2005) *Raphignathoidea* (Acari: Prostigmata). *Fauna of New Zealand* 52. Mannaki Whenua Press, Lincoln, New Zealand, 400 pp.
- Grandjean F (1939) Les segments postlarvaires de l'hysterosoma chez les oribates (acariens). *Bulletin de la Société Zoologique de France* 64: 273–284.
- Grandjean F (1944) Observation sur les acariens de la famille des Stigmaeidae. *Archives des Sciences Physiques et Naturelles* 26: 103–131.
- Grandjean F (1946) Au sujet de l'organe de Claparède, des eupathides multiples et des taenidies mandibulaires chez les Acariens actinochitineux. *Extrait des Archives des Sciences Physiques et Naturelles* 28: 63–87.
- Kethley J (1990) Acarina: Prostigmata (Actinedida). In: Dindal DL (Ed.) *Soil Biology Guide*. John Wiley & Sons, New York, 667–756.
- Khanjani M, Pishavar S, Mirmoayed A (2013) A new species of *Raphignathus* Dugés (Acari: Raphignathidae) from Iran. *Acarina (Moskva)* 21: 62–68.
- Kramer P (1877) Grundzüge zur Systematik der Milben. *Archiv für Naturgeschichte* 43(1): 215–247.
- Krantz GW, Walter DE (2009) *A Manual of Acarology*. Texas Tech University Press, Lubbock, Texas, 806 pp.
- Meyer MKP, Ryke PAJ (1960) Mites of the superfamily Raphignathoidea (Acarina: Prostigmata) associated with South African plants. *Annales and Magazine of Natural History (Series 13)* 2: 209–234. <https://doi.org/10.1080/00222935908651024>
- Meyer MKPS, Ueckermann EA (1989) *African Raphignathoidea* (Acari: Prostigmata). *Entomology Memoir, Department of Agricultural Technical Services. Republic of South Africa* 74: 1–58.

- Nasrollahi S, Khanjani M, Mirfakhraee S (2018) A new species of the genus *Raphignathus* (Acari: Raphignathidae) from Kurdistan, Iran, with a key to world species. *Systematic and Applied Acarology* 23(10): 2070–2081. <https://doi.org/10.11158/saa.23.10.14>
- Pishevvar SH, Khanjani M (2021) Two new species of the genus *Raphignathus* (Acari: Raphignathidae) from Hamadan, Iran, with a key to the known species. *Systematic and Applied Acarology* 26(1): 185–198. <https://doi.org/10.11158/saa.26.1.11>
- Rack G (1962) Milben aus Taubennestern mit Beschreibung einer neuen Art, *Acheles gracilis* (Acarina, Raphignathidae). *Zoologischer Anzeiger* 168(7–10): 275–292.
- Smiley RL, Moser JC (1968) New species of mites from pine (Acarina: Tarsocheylidae, Eupalopsellidae, Caligonellidae, Cryptognathidae, Raphignathidae and Neophyllobidae). *Proceedings of the Entomological Society of Washington* 70(40): 307–317.